

# Some Results on Derivatives of Generalized Modified Baskakov Type Operators (A)

Dr. R.P. Pathak

Department of Mathematics,  
National Institute of Technology, Raipur  
G.E. Road, Raipur -492010 (C.G.) INDIA

Shiv Kumar Sahoo

Department of Mathematics,  
National Institute of Technology, Raipur  
G.E. Road, Raipur -492010 (C.G.) INDIA

Ashish Kumar Tiwari

Department of Mathematics,  
National Institute of Technology, Raipur  
G.E. Road, Raipur -492010 (C.G.) INDIA

**Abstract** – Recently Deo N.et.al. (Appl. Maths. Compt., 201(2008), 604-612.) introduced a new Bernstein type special operators. Motivated by Deo N.et.al., in this paper we introduce generalization of positive linear operators (4) which is generator of positive linear operators (6) and (7). We shall study some approximation results on it.

**Keywords** – Positive linear operators, Bernstein type special operators, Baskakov type positive linear operators, Sz sz type positive linear operators.

## I. INTRODUCTION

Baskakov [6] introduced the sequences of positive linear operators  $\{L_n\}$ ,  $(n \in \mathbb{N})$   $L_n \subset [0, \infty) \subset [0, R]$   $(R > 0)$  which are defined by

$$(L_n f)(x) = \sum_{k=0}^{\infty} \frac{(-x)^k}{k!} \phi_n^{(k)}(x) f\left(\frac{k}{n}\right) \dots (1)$$

and are generated by sequence of functions  $\{\phi_n\}$ ,  $(n \in \mathbb{N})$  possesses the following five properties :

- (a)  $\phi_n$  is analytic on the interval  $[0, R]$  where  $R > 0$  including end point.
- (b)  $\phi_n(0) = 1$ .
- (c)  $\phi_n$  is completely monotone i.e.  $(-1)^k \phi_n^{(k)}(x) \geq 0$  if  $k = 0, 1, 2, \dots$
- (d) There exist positive integer  $m(n)$  not depending on  $k$  such that

$$-\phi_n^{(k)}(x) = n \phi_{m(n)}^{(k-1)}(x) [1 + \alpha_{k,n}(x)]$$

where  $\alpha_{k,n}(x)$  converges to zero uniformly in  $k$  as  $n \rightarrow \infty$ .

- (e)  $\lim_{n \rightarrow \infty} \frac{n}{m(n)} = 1$ .

Deo N.et.al. [1] introduced a new Bernstein type special operators  $\{V_n f\}$  defined as,

$$(V_n f)(x) = \sum_{k=0}^n p_{n,k}(x) f\left(\frac{k}{n}\right) \dots (2)$$

where

$$p_{n,k}(x) = \left(1 + \frac{1}{n}\right)^n \binom{n}{k} x^k \left(\frac{n}{n+1} - x\right)^{n-k};$$

for  $0 \leq x \leq \frac{n}{n+1}$

Again Deo N.et.al. [1] gave the integral modification of the operators (2) which are defined as,

$$(L_n f)(x) = n \left(1 + \frac{1}{n}\right)^n \times \sum_{k=0}^n p_{n,k}(x) \int_0^{\frac{n}{n+1}} p_{n,k}(t) f(t) dt \dots (3)$$

and prove some approximation results on the operators (3).

Motivated by Deo N.et.al.[1] we studied a sequence of positive linear operators  $\{M_n f\}$  which are defined as,

$$(M_n f)(x) = n \left(1 + \frac{1}{n}\right)^2 \times \sum_{k=0}^n \frac{(-x)^k}{k!} \phi_n^{(k)}(x) \int_0^{\frac{n}{n+1}} p_{n,k}(t) f(t) dt \dots (4)$$

where

$$p_{n,k}(t) = \left(1 + \frac{1}{n}\right)^n \binom{n}{k} t^k \left(\frac{n}{n+1} - t\right)^{n-k};$$

$p > 0$  and for  $t \in \left[0, \frac{n}{n+1}\right]$ .

and are generated by sequence of functions  $\{\phi_n\}$ ,  $(n \in \mathbb{N})$  possesses the above mentioned five properties. We shall study some approximation results on the operators (4).

The following are some special cases of the operators (4) :

- A.  $\phi_n(x) = e^{-nx}$
- B.  $\phi_n(x) = (1-x)^n$
- C.  $\phi_n(x) = (1+x)^{-n}$ .

In this paper we shall study some results on the positive linear operator (4) for special case  $\phi_n(x) = e^{-nx}$ .

## II. MAIN RESULTS

In order to complete our study first we show that  $\phi_n(x) = e^{-nx}$  satisfies the above five properties (a) to (e).

- (a) Let  $\phi_n(x) = e^{-nx}$

Now successive differentiate the  $\phi_n(x)$ ,  $k$  times w.r.t.  $x$ , we get

$$\begin{aligned} \phi_n'(x) &= -n e^{-nx} \\ \phi_n''(x) &= (-1)^2 n^2 e^{-nx} \\ \phi_n'''(x) &= (-1)^3 n^3 e^{-nx} \end{aligned}$$

.....

$$\phi_n^{(k)}(x) = (-1)^k n^k e^{-nx} \dots (5)$$

Therefore  $\phi_n(x)$  is differentiable in the interval  $\left[0, \frac{n}{n+1}\right]$ .

- (b)  $\phi_n(0) = e^{-n(0)} = e^0 = 1$ .

- (c) If  $\phi_n(x) = e^{-nx}$

$$\begin{aligned} \phi_n^{(k)}(x) &= (-1)^k n^k e^{-nx} \\ (-1)^k \phi_n^{(k)}(x) &= (-1)^k (-1)^k n^k e^{-nx} \\ &= (-1)^{2k} \frac{n^k}{e^{nx}} \geq 0. \end{aligned}$$

- (d) There exist positive integer  $m(n)$  not depending on  $k$  such that

$$-\phi_n^{(k)}(x) = n \phi_{m(n)}^{(k-1)}(x) \left|1 + \lim_{n \rightarrow \infty} \alpha_{k,n}(x)\right|$$

where  $\lim_{n \rightarrow \infty} \alpha_{k,n}(x) = 0$ .  
 since  $\phi_n^{(k)}(x) = (-1)^k n^k e^{-nx}$   
 $\phi_n^{(k-1)}(x) = (-1)^{k-1} n^{k-1} e^{-nx}$   
 therefore  
 $(-1) \phi_n^{(k)}(x) = (-1)(-1)^k n^k e^{-nx}$   
 $- \phi_n^{(k)}(x) = (-1)^{k-1} n^{k-1} e^{-nx}$   
 $= n \phi_n^{(k-1)}(x)$   
 $= n \phi_n^{(k-1)}(x) \left[ 1 + \lim_{n \rightarrow \infty} \alpha_{k,n}(x) \right]$ .

(e)  $\lim_n \frac{n}{m(n)} = \lim_n \frac{n}{n+c} = 1$ .

Here we took  $c = 0$ .

This completes the proof.

If we put the value of  $\phi_n^{(k)}(x)$  in the operator (4), we get new modified positive linear operator.

$$\begin{aligned} (M_n f)(x) &= n \left( 1 + \frac{1}{n} \right)^2 \times \\ &\sum_{k=0}^{\infty} \frac{(-x)^k}{k!} (-1)^k n^k e^{-nx} \int_0^n p_{n,k}(t) f(t) dt \\ &= n \left( 1 + \frac{1}{n} \right)^2 \times \\ &\sum_{k=0}^{\infty} \frac{(-1)^{2k} (nx)^k}{k!} e^{-nx} \int_0^n p_{n,k}(t) f(t) dt \\ &= n \left( 1 + \frac{1}{n} \right)^2 \times \\ &e^{-nx} \sum_{k=0}^{\infty} \frac{(nx)^k}{k!} \int_0^n p_{n,k}(t) f(t) dt \quad \dots (6) \end{aligned}$$

Which is modified Sz sz type positive linear operators and it is given by us.

Now we are modifying operator (6) for  $p > 0$  as follows:

$$\begin{aligned} (M_n f)(x) &= n \left( 1 + \frac{1}{n} \right)^2 e^{-(n+p)x} \\ &\sum_{k=0}^{\infty} \frac{(n+p)^k (x)^k}{k!} \int_0^n p_{n,k}(t) f(t) dt \quad \dots (7) \end{aligned}$$

Which is modified Szasz type positive linear operators and it is given by us.

### III. CONCLUSION

In this paper we conclude that our new modified Baskakov type operators (4) is generalized positive linear operator of modified Sz sz type operators (6) & (7). These operators have different approx properties, analysis is different. We will discuss them elsewhere.

### IV. ACKNOWLEDGMENT

The authors are thankful to Director of National Institute of Technology, Raipur (C.G.) for encouragement.

### REFERENCES

- [1] Deo N., Noor M.A. and Siddiqui M.A. "On approximation by class of new Bernstein type operators", Appl. Math. Comput., 201 (2008), 604-612.
- [2] Kasana H.S. , Prasad G., Agrawal P.N. and Sahai A. "Modified Sz sz operators" Proceeding of International Conference on Mathematical Analysis and its Applications Pergamon Press, 1985, 29-41.
- [3] Papanicolaou C.G. "Some Bernstein type operators" Amer. Math. Month, 82 (1975) ,674-677.
- [4] Singh S.P., "Some problems on approximation of functions by positive linear operators" Ph.D. Thesis, Roorkee University (1982).
- [5] Shiv Kumar Sahoo and S. P. Singh "Some approximation results on a special class of positive linear operators" , Proceedings of The Mathematical Society , Banaras Hindu University, Vol. 24 (2008), 1-9.
- [6] Baskakov V. A., "An example of a sequence of linear positive operators in the space of continuous functions", Dokl. Akad. Nauk, 13(1973), 249-251.

### AUTHOR'S PROFILE

#### Dr. R.P. Pathak

Associate Professor  
 Department of Mathematics,  
 National Institute of Technology, Raipur  
 G.E. Road, Raipur -492010 (C.G.)  
 E-mail id : [r.p.pathak8@gmail.com](mailto:r.p.pathak8@gmail.com)

#### Shiv Kumar Sahoo

Department of Mathematics,  
 National Institute of Technology, Raipur  
 G.E. Road, Raipur -492010 (C.G.)  
 E-mail id: [maths10sks@gmail.com](mailto:maths10sks@gmail.com).

#### Ashish Kumar Tiwari

Department of Mathematics,  
 National Institute of Technology, Raipur  
 G.E. Road, Raipur -492010 (C.G.)  
 E-mail id : [ashish0584@gmail.com](mailto:ashish0584@gmail.com).